

AMOXYLIFE ADVANCE®

SYNERGY IN ACTION

Amoxicillin
150 mg

AMOXYLIFE
ADVANCE®

Amoxicillin 150 mg
Gentamicin 40 mg

Suspension for Injection

Sterile

For S.C./I.M. Use Only



STORAGE CONDITIONS
Do not store above 25°C. Keep this carton in order to protect from light.
Discard unused material safely.
Keep away from children.

MARKETING AUTHORISATION NO:
WORLD-VET İLAÇ SAN. VE TİC. LTD. ŞTİ.
15 Temmuz Mah. Cami Yolu Cad. No:
K1 Göneşli, Bağcılar, İstanbul 34290



Gentamicin
40 mg

AMOXYLIFE ADVANCE®

SYNERGY IN ACTION

ADVANTAGES

- Broad Spectrum of Activity
- Less Resistance
- Fewer Side Effects
- Easy to inject

INDICATIONS

Gastrointestinal and urogenital infections, respiratory and soft tissue infections, postoperative infections, pneumonia, enteritis, and peritonitis. Supporting therapy in addition to local treatment of mastitis and metritis.



Bacterial infections caused by microorganisms susceptible to amoxicillin and gentamicin.

| PATOGENS | <i>Escherichia coli</i> | <i>Citrobacter spp.</i> | <i>Klebsiella spp.</i> | <i>Pasteurella spp.</i> | <i>Proteus spp.</i> | <i>Salmonella spp.</i> | <i>Actinomyces pyogenes</i> | <i>Staphylococcus spp.</i> | <i>Streptococcus spp.</i> | <i>Bacillus spp.</i> | <i>Corynebacterium spp.</i> | <i>Clostridium spp.</i> | <i>Bacteroides spp.</i> | <i>Fusobacterium spp.</i> | <i>Actinomycetilus spp.</i> | <i>Erysipelothrix rhusiopathiae</i> |
|--------------------|-------------------------|-------------------------|------------------------|-------------------------|---------------------|------------------------|-----------------------------|----------------------------|---------------------------|----------------------|-----------------------------|-------------------------|-------------------------|---------------------------|-----------------------------|-------------------------------------|
| AMOXICILLIN | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● |
| GENTAMICIN | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● |
| AMOXYLIFE ADVANCE® | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● |

● Partially active ● Active ● Highly active

DOSAGE AND ADMINISTRATION

Cattle, sheep and goats, pigs: 1 ml/10 kg BW, i.m.

Cats & dogs: 1 ml/10 kg BW, i.m. or s.c.