

# PRODUCT GUIDE

	TRADE NAME	ACTIVE SUBSTANCES	DOSAGE AND ADMINISTRATION	PRINCIPAL INDICATIONS	TARGET SPECIES
<b>ANTIBIOTICS</b>					
1	AMOXYLIFE-LA® Suspension for Injection	Amoxicillin 150 mg	Cattle, sheep and goats, pigs: 1 ml/10 kg BW, i.m. Cats & dogs: 1 ml/10 kg BW, i.m. or s.c.	Gastrointestinal and urogenital infections, respiratory and soft tissue infections, postoperative infections, pneumonia, enteritis, and peritonitis. Supporting therapy in addition to local treatment of mastitis and metritis	
2	AMOXYLIFE ADVANCE® Suspension for Injection	Amoxicillin 150 mg Gentamicin 40 mg	Cattle, sheep and goats, pigs: 1 ml/10 kg BW, i.m. Cats & dogs: 1 ml/10 kg BW, i.m. or s.c.	Bacterial infections caused by microorganisms susceptible to amoxicillin and gentamicin	
3	AZITROSTAR® Solution for Injection	Azithromycin 100 mg	Cattle, sheep and pigs: 1 ml/20 kg BW, i.m. once daily	Infections of the respiratory system, digestive and genitourinary system, infections of skin and soft tissues, necrobacteriosis; erysipelas of pigs; spirochetosis; mycoplasma infections; conjunctivitis	
4	CEFQUITAL® Suspension for Injection	Cefquinome 25 mg	Cattle, sheep and goats: 1 ml/25 kg BW i.m. Pigs: 1-2 ml/25 kg BW i.m. Sows: 2 ml/25 kg BW i.m. Cats & dogs: 0.5 ml/5 kg BW i.m. or s.c.	Respiratory tract infections and foot diseases caused by cefquinome susceptible bacteria, <i>E.coli</i> derived calf septicemias, acute mastitis infections, and secondary infections associated with viral diseases	
5	CEFTIVET® Suspension for Injection	Ceftiofur 50 mg	Cattle, sheep and goats: 1 ml/50 kg BW i.m. Pigs: 1 ml/16 kg BW i.m. Horses: 1 ml/50 kg BW i.m. Dogs: 0.2 ml/5 kg BW i.m.	BRD, acute bovine interdigital necrobacillosis, and acute metritis associated with microorganisms susceptible to ceftiofur	
6	CEFTIVET® ADVANCE Suspension for Injection	Ceftiofur 50 mg Ketoprofen 150 mg	Cattle: 1 ml/50 kg BW i.m.	Treatment of acute and chronic laminitis, acute infectious mastitis, dystocia, and respiratory diseases with inflammatory reaction associated with microorganisms susceptible to ceftiofur	
7	EVERODOX WSP® Oral solution	Doxycycline 500 mg	Cattle, sheep: 3-4 mg/kg twice daily Pigs: 5-8 mg/kg twice daily Poultry: 0.2 ml/L of drinking water for 3-5 days	Colibacteriosis, salmonellosis, staphylococcosis, pasteurellosis, hemophilia, swine enzootic pneumonia, swine erysipelas, hoof rot, keratoconjunctivitis, arthritis, abscess, and other diseases caused by bacteria and mycoplasma	
8	ENFLOREX® Solution for Injection	Enrofloxacin 100 mg	Cattle: 0.5 ml/10 kg BW i.m. Sheep and goats: 0.5 ml/20 kg BW i.m. Pigs: 1 ml/40 kg BW i.m. Dogs: 0.5 ml/10 kg BW i.m.	Gastrointestinal and respiratory infections caused by enrofloxacin-sensitive microorganisms	
9	ENFLOREX® Oral Solution	Enrofloxacin 100 mg	Cattle, sheep and goats, pigs: 1 ml/20 kg BW daily for 3-5 days Poultry: 1 ml/20 kg of live weight daily, added into drinking water	Colibacteriosis, salmonellosis, streptococcosis, mycoplasmosis, necrotic enteritis, and other infections associated with enrofloxacin-sensitive microorganisms	
10	FLORRELIEF® Solution for Injection	Florfenicol 300 mg Flunixin 16,5 mg	Cattle: 2 ml/15 kg BW s.c. in the neck	Treatment of BRD associated with <i>Mannheimia haemolytica</i> , <i>Pasteurella multocida</i> , <i>Histophilus somni</i> , and <i>Mycoplasma bovis</i> . Control of BRD-associated pyrexia in beef and non-lactating dairy cattle	
11	LUFLOLAN® Solution for Injection	Florfenicol 300 mg	Cattle, sheep and goats: 1 ml/15 kg BW i.m. in the neck twice with a 48-hour interval between doses or 2 ml/15 kg BW s.c. as a single injection Pigs: 1 ml/20 kg BW i.m. in the neck twice with a 48-hour interval	Respiratory infections, limb infections, bacterial diarrhoea in calves	
12	LEVOVITAL® Solution for Injection	Levofloxacin 150 mg	Cattle, swine: 1 ml/30 kg BW i.m. for 3-5 days	Infections associated with levofloxacin-sensitive microorganisms	
13	LEVOVITAL® Oral Solution	Levofloxacin 150 mg	Swine, poultry: 1 ml/20 kg BW for 3-5 days	Infections associated with levofloxacin-sensitive microorganisms	
14	MARBOPRIME® Solution for Injection	Marbofloxacin 100 mg	Cattle: 2 ml/25 kg BW for the treatment of respiratory infections 1 ml/50 kg BW for the treatment of mastitis Pigs: 1 ml/50 kg BW i.m. in the neck	Respiratory infections, mastitis, MMA syndrome	
15	SPIRALIFE® Solution for Injection	Spiramycin 600 000 EU	Cattle: 5 ml/100 kg BW Calves, pigs: 5 ml/40 kg BW i.m.	Gastrointestinal and respiratory infections caused by spiramycin-sensitive microorganisms	
16	TETRALIFE-LA® Solution for Injection	Oxytetracycline 200 mg	Cattle, sheep and goats, pigs: 1 ml/10 kg BW, i.m. once in 3 days, but not more than 3 times	Infections associated with oxytetracycline-sensitive microorganisms	
17	TETRALIFE ADVANCE® Solution for Injection	Oxytetracycline 300 mg Flunixin 20 mg	Cattle: 1 ml/10 kg BW i.m. as a single injection	Infections associated with oxytetracycline-sensitive microorganisms	
18	TULATRAM® Solution for Injection	Tulathromycin 100 mg	Cattle: 1 ml/40 kg BW s.c. as a single injection Pigs: 1 ml/40 kg BW i.m. as a single injection in the neck	Treatment of BRD associated with <i>M. haemolytica</i> , <i>P. multocida</i> , <i>H. somni</i> , and <i>M. bovis</i> Treatment of SRD associated with <i>A. pleuropneumoniae</i> , <i>P. multocida</i> , <i>B. bronchiseptica</i> , <i>H. parasuis</i> , and <i>M. hyopneumoniae</i> ; control of SRD associated with <i>A. pleuropneumoniae</i> , <i>P. multocida</i> , and <i>M. hyopneumoniae</i>	
<b>INTRAMAMMARY TUBES</b>					
19	COMBIMAST® Intramammary Suspension (for lactating cattle)	Each syringe (5 g) contains: Procaine penicillin G 100.000 IU Streptomycin sulphate 100 mg Neomycin sulphate 100 mg Prednisolone 10 mg	1 syringe into each affected quarter at 24-hour intervals for 3 consecutive milkings	For use in clinical cases of mastitis	
20	NAFCIVET-DC® Intramammary Suspension (dry period)	Each syringe (3 g) contains: Procaine benzylpenicillin G 300 mg Dihydrostreptomycin 100 mg Nafcilin 100 mg	1 syringe into each affected quarter after lactation period	Prevention of mastitis in dry cows	
<b>ANTIPARASITICS</b>					
21	DIFFICARB® Solution for Injection	Imidocarb 85 mg	Cattle, sheep and goats: 0.1 ml/10 kg i.m. Horses: 2 ml/100 kg i.m. Dogs: 0.25-0.5 ml/10 kg s.c.	Prevention of babesiosis and anaplasmosis	
22	IVERVEXAN® Solution for Injection	Ivermectin 10 mg	Cattle, horses: 1 ml/50 kg BW s.c. Sheep and goats: 0.5 ml/25 kg BW s.c. Pigs: 1 ml/33 kg BW s.c. Rabbits: 0.1 ml/5 kg BW	Treatment and control of gastrointestinal roundworms, lungworms, grubs, sucking lice, chorioptic and sarcoptic mange mites	
23	DORAVEXAN® Solution for Injection	Doramectin 10 mg	Cattle: 1 ml/50 kg BW s.c. Sheep and goats: 1 ml/50 kg BW i.m. Pigs: 1 ml/33 kg BW i.m.	Treatment and control of gastrointestinal roundworms, lungworms, eyeworms, grubs, sucking lice, and mange mites	
24	EPRINYSE® Solution for Injection	Eprinomectin 10 mg	Cattle, sheep and goats, horses: 1 ml/50 kg BW s.c. or i.m. Pigs: 1 ml/33 kg BW i.m. Cats & dogs: 0.2 ml/10 kg BW	Treatment and control of internal and external parasites of beef and dairy cattle, including lactating dairy cows	
25	NAMEDOL® Oral Suspension	Albendazole 100 mg	Cattle: 0.75-1 ml/10 kg BW Sheep and goats: 0.75-1.5 ml/10 kg BW	Prevention and treatment of worm infections	
<b>ANTICOCCIDIALS</b>					
26	TOLTRAZOR® 2,5% Oral Solution	Toltrazuril 25 mg	28 ml/100 kg of live weight daily, added into drinking water	Prevention and treatment of eimeriosis (coccidiosis)	
27	TOLTRAZOR® 5% Oral Suspension	Toltrazuril 50 mg	Calves: 3 ml/10 kg BW Lambs: 0.4 ml/kg BW Piglets: 1 ml/2.5 kg BW	Prevention and treatment of eimeriosis (coccidiosis)	
<b>ANTI-INFLAMMATORY AGENTS</b>					
28	DEXAPRIME® Solution for Injection	Dexamethasone 4 mg	Horses, cattle: 5 ml Pigs: 2-3 ml Piglets, dogs: 0.5-1 ml Cats: 0.25-0.5 ml once daily for 3 days	Allergic reactions, myoglobinuria, toxinaemia, shock, stress and urticaria. Local treatment of arthritis, bursitis, distortions, peri-arthritis, tendinitis and tendovaginitis	
29	FLUNALGIN® Solution for Injection	Flunixin 50 mg	Horses: 1 ml/45 kg s.c. or i.v. Cattle, pigs: 2 ml/45 kg s.c. or i.v.	Anti-inflammatory, antipyretic and analgesic agent	
30	KETAPRIN® Solution for Injection	Ketoprofen 100 mg	Cattle: 3 ml/100 kg BW i.v. or i.m. Pigs: 3 ml/100 kg BW i.m. Horses: 1 ml/45 kg BW i.m.	Anti-inflammatory, antipyretic and analgesic agent	
31	LOMECAM® Solution for Injection	Meloxicam 20 mg	Cattle: 2.5 ml/100 kg BW s.c. or i.v. Pigs: 2.0 ml/100 kg BW i.m.	Anti-inflammatory, antipyretic and analgesic agent	
<b>VITAMINS AND MINERALS</b>					
32	SELENVET-B® Emulsion for Injection	Sodium selenite 1 mg Vitamin E 60 mg Vitamin B1 40 mg	Cattle, sheep and goats: 1 ml/50 kg BW s.c. or i.m.	Prevention and therapy of diseases associated with deficiency of vitamin E and selenium, especially muscular dystrophy in young livestock; beneficial effect on the reproduction of cows	
33	VITABUTAN® Solution for Injection	Butaphosphan 100 mg Vitamin B12 0.05 mg	Cattle, horses: 5-25 ml per animal Sheep and goats: 2.5-8 ml per animal Pigs: 2.5-10 ml per animal Poultry: 1-3 ml/L of drinking water	Prevention or treatment of deficiencies of Vitamin B12 and phosphorus in cattle, swine, horses, and poultry	
34	BETACARTIN-AQUA® Solution for Injection	Beta-carotene 10 mg	Horse: 20 ml for 1-3 days for 14 day i.m. Cattle: 20-25 ml for 14 days i.m. or s.c. Calves: 5 ml immediately after birth i.m. or s.c. Pigs: 7 ml on the day of insemination	Metabolism-boosting supplement used for improvement of performance	
35	MULTIWWET® Solution for Injection	Each ml contains: vitamin A 50 000 EU; vitamin D3 25 000 EU; vitamin E 4 mg; vitamin B1 10 mg; vitamin B2 0.04 mg; vitamin B6 1 mg; vitamin B12 0.01 mg; vitamin B5 2 mg; vitamin B3 5 mg; vitamin B8 2 mg; methionine 5 mg; choline citrate 5 mg; magnesium hypophosphite 1 mg; cobalt chloride 0.02 mg; copper sulfate 0.1 mg; zinc sulfate 0.1 mg; manganese sulfate 0.1 mg; selenium 0.7 mg; benzyl alcohol 15 mg; water for injection q.s. to 1 ml	Cattle, horses: 2-6 ml/100 kg BW Pigs, sheep and goats: 5-8 ml/50 kg BW Piglets, calves, kids and lambs: 1-1.5 ml/10 kg BW Dogs, cats: 0.5-1 ml/10 kg BW	Prevention and treatment of vitamin deficiencies, growth disturbances, weakness of newborn animals, neonatal anemia, visual impairment, gastrointestinal disorders, anorexia, non-infectious reproductive disorders, rachitis, muscle weakness, muscle tremor, and myocardial failure with breath disorders	